

TGFB1/ TGFβ1/ TGF-β1, Human, Recombinant

货号 : PCK091

产品信息

| | |
|------|--|
| 别名 | Transforming Growth Factor Beta-1; TGF-Beta-1; Latency-Associated Peptide; LAP; TGFB1; TGFB |
| 物种 | Human |
| 表达宿主 | CHO Stable Cells |
| 序列信息 | Ala279-Ser390 |
| 检索号 | P01137 |
| 分子量 | 12.8 kDa |
| 生物活性 | Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is 4-40 pg/ml. |

产品特性

| | |
|-----|---|
| 纯度 | >95% as determined by reducing SDS-PAGE. |
| 内毒素 | <1.0 EU per µg as determined by LAL test. |
| 保存 | Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months. |
| 运输 | Ambient temperature or ice pack. |
| 制剂 | Lyophilized from a 0.2 µm filtered solution of 50mM Glycine-HCl, 150mM NaCl, pH2.5. |




复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


背景介绍

Transforming Growth Factor β -1 (TGF β -1) is a secreted protein which belongs to the TGF- β family. TGF β -1 is abundantly expressed in bone, articular cartilage and chondrocytes and is increased in osteoarthritis (OA). TGF β -1 performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation and apoptosis. The precursor is cleaved into a latency-associated peptide (LAP) and a mature TGF β -1 peptide. TGF β -1 may also form heterodimers with other TGF β family members. It has been found that TGF β -1 is frequently upregulated in tumor cells. Mutations in this gene results in Camurati-Engelmann disease.

SDS-PAGE



生物活性



Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 human erythroleukemic cells. The ED50 for this effect is 4-40 pg/ml.

