

#### ACVR2A (C-6His), Human, Recombinant

货号: PCK106

### 产品信息

- 别名 Activin Receptor Type-2A; Activin Receptor Type IIA; ACTR-IIA; ACTRIIA; Pricella ACVR2A; ACVR2
- 物种 Human
- Human Cells 表达宿主
- 序列信息 Ala20-Pro134
- 检索号 P27037
- 分子量 14.35 kDa

#### 产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	< 1.0 EU per μg as determined by LAL test.
保存	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
运输	Ambient temperature or ice pack.
制剂	Supplied as a 0.2 µm filtered solution of 20mM PB, 300mM NaCl, 10% Glycerol, 5% Trealose, pH7.4.






## 背景介绍

**SDS-PAGE** 

Activin Receptor Type-2A is a Protein that in humans is encoded by the ACVR2A gene. ACVR2A is an activintype 2 Receptor. This gene encodes activin A type II Receptor. Activins are dimeric growth anddifferentiation factors which belong to the transforming Growth Factor-beta (TGF-beta) superfamily ofstructurally related signaling Proteins. Activins signal through a heteromeric complex of Receptor serinekinases which include at least two type I (I and IB) and two type II (II and IIB) Receptors. These Receptors areall transmembrane Proteins, composed of a Ligand-binding extracellular domain with cysteine-rich region, atransmembrane domain, and a cytoplasmic domain with predicted serine/ threonine specificity. Type IReceptors are essential for signaling; and type II Receptors form a stable complex after Ligand binding, resultingin phosphorylation of type I Receptors by type II Receptors. Type II Receptors are considered to beconstitutively active kinases.



# ma pricella

