

FGF-17 (C-6His), Human, Recombinant

货号: PCK130

产品信息

别名 Fibroblast Growth Factor 17; FGF-17; FGF17 Pricella

物种 Human

表达宿主 **Human Cells**

序列信息 Thr23-Thr216

检索号 O60258

分子量 22.6 kDa

标签 C-6His

产品特性

纯度 >95% as determined by reducing SDS-PAGE.

内毒素 < 1.0 EU per µg as determined by LAL test.

保存 Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at -5~-20°C for 3 months.

运输 Ambient temperature or ice pack.

制剂 Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋








复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml.Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Fibroblast Growth Factor 17 (FGF17) is a member of the heparin-binding Growth Factors family that is prominently expressed in the cerebellum and cortex. Proteins of this family possess broad mitogenic and cell survival activities and they are involved in a variety of biological processes including embryonic development cell growth, morphogenesis, tissue repair, tumor growth, and invasion. FGF17 plays an important role in the regulation of embryonic development and it acts as signaling molecule in the induction and patterning of the embryonic brain. In addition, FGF17 stimulates the proliferation and activation of cells that express FGF Receptors.

SDS-PAGE



网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



