

IL-33/IL-1F11, Human, Recombinant

货号 : PCK191

产品信息

别名	IL-1 F11, NF-HEV, DVS27
物种	Human
表达宿主	E.coli
序列信息	MSITGISPITEYLASLSTYNDQSITFALEDESYEIYVEDLKKDEKKDKVLLSYY ESQHPSNESGDGVDGKMLMVTLSPTKDFWLHANNKEHSVELHKCEKPLPD QAFFVLHNMHSNCVSFECKTDPGVFIGVDNHLALIKVDSSENLCTENILFK LSET with polyhistidine tag at the C-terminus.
检索号	O95760.1
分子量	18.93 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to binding with recombinant ST2L (IL-33 receptor). The ED50 for this effect is <54 ng/mL. Measure by its ability to induce proliferation in D10.G4.1 cells. The ED50 for this effect is < 0.1 ng/mL. The specific activity of recombinant human IL-33 is approximately >4 x10 ⁷ IU/ mg.

产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin 33 (IL-33) is a member of the IL-1 family that potently drives production of T helper-2 (Th2)-associated cytokines (e.g., IL-4). IL33 is a ligand for ST2 (IL1RL1), an IL-1 family receptor that is highly expressed on Th2 cells, mast cells and group 2 innate lymphocytes. It is expressed by a wide variety of cell types, including fibroblasts, mast cells, dendritic cells, macrophages, osteoblasts, endothelial cells, and epithelial cells.

SDS-PAGE

