

IL-4Rα/CD124 (C-mFc), Human, Recombinant

货号: PCK198

产品信息

别名 Interleukin-4 Receptor subunit alpha; IL-4 Receptor subunit alpha; IL-4R subunit

alpha; IL-4R-alpha; IL-4RA; CD124; IL-4-binding Protein; IL4-BP; IL4R; IL4RA

物种 Human

表达宿主 Human Cells ?

序列信息 Met26-Gln231

检索号 P24394

分子量 50.2 kDa

生物活性 Loaded Human IL-4-His on HIS1K Biosensor, can bind Human IL-4RA-mFc with

an affinity constant of 0.12 nM as determined in BLI assay.

产品特性

纯度 >95% as determined by reducing SDS-PAGE.

内毒素 < 1.0 EU per µg as determined by LAL test.

保存 Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at -5~-20°C for 3 months.

运输 Ambient temperature or ice pack.

制剂 Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋








复融


Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

Interleukin 4 Receptor alpha (IL4-Ra) is a widely expressed 140 kDa transmembrane glyco Protein in the class I Cytokine Receptor family. Mature human IL4-Ra consists of a 207 amino acid (aa) extracellulardomain (ECD) that contains a Cytokine binding region and one fibronectin type III domain, a 24 aatransmembrane segment, and a 569 aa cytoplasmic domain that contains one Box 1 Motif and one ITIMMotif. IL4-Ra plays an important role in Th2-biased immune responses, alternative macrophage activation, mucosal immunity, allergic inflammation, tumor progression, and atherogenesis. Soluble forms of IL4-Ra, generated by alternate splicing or proteolysis, retain Ligand binding properties and inhibit IL-4 bioactivity.

SDS-PAGE





Loaded Human IL-4-His on HIS1K Biosensor, can bind Human IL-4RA-mFc with an affinity constant of 0.12 nM as determined in BLI assay

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



