

## VEGF/ VEGF-A, Rat, Recombinant

货号 : PCK266

### 产品信息

别名	Vascular endothelial Growth Factor A; Vascular permeability factor; VEGF; VEGF-A; VPF
物种	Rat
表达宿主	Yeast
序列信息	Ala27-Arg190(Ala36Thr)
检索号	P16612-2
分子量	19.2 kDa
生物活性	Immobilized Rat VEGF 164 at 2µg/ml (100 µl/well) can bind Mouse VEGFR2-Fc. The ED50 of Mouse VEGFR2-Fc is 22.17 ng/ml.

### 产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	< 1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.




## 复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


## 背景介绍

Vascular endothelial Growth Factor (VEGF/ VEGF-A) is originally known as vascular permeability factor (VPF). It belongs to the PDGF family with a cysteine-knot structure comprised of eight conserved cysteine residues, and reckoned as a potent mediator in the process of angiogenesis and vasculogenesis in either fetus or adult. VEGF is particularly expressed in supraoptic, paraventricular nuclei and the choroid plexus of the pituitary, and abundant in the corpus luteum of the ovary and in kidney glomeruli. The rat VEGF Protein contains a putative 20 amino acids (aa) signal peptide, and alternative splicing of rat VEGF gene produces isoforms of 120, 144, 164 and 188 aa. Rat VEGF164 respectively displays 97% and 88% aa identity with that regions of mouse and human VEGF. VEGF can bind to the FLT1/ VEGFR1 and KDR/ VEGFR2 Receptors, heparan sulfate and heparin, and play important roles in inducing endothelial cell proliferation, promoting cell migration, inhibiting apoptosis and inducing permeabilization of blood vessels.

## SDS-PAGE



## 生物活性



Immobilized Rat VEGF 164 at 2 µg/ml (100 µl/well) can bind Mouse VEGFR2-Fc. The ED50 of Mouse VEGFR2-Fc is 22.17 ng/ml.

