

IL-3RA, Human, Recombinant

货号：PCK075

产品信息

别名	Interleukin-3 Receptor subunit alpha; IL-3 Receptor subunit alpha; IL-3R subunit alpha; IL-3R-alpha; IL-3RA
物种	Human
表达宿主	Human Cells
序列信息	Thr19-Arg305
检索号	P26951
分子量	60 kDa

产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	< 1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.



复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

背景介绍

CD123, also known as Interleukin-3 Receptor subunit alpha, belongs to the type I Cytokine Receptor family. In mouse, there are two classes of high-affinity IL3 Receptors. One contains an IL3-specific beta subunit and the other contains the beta subunit also shared by high-affinity IL5 and GM-CSF Receptors. CD123 stimulates the proliferation and differentiation of hemopoietic cells including the pluripotent hematopoietic stem cells as well as various lineage-committed cells. CD123 is a heterodimer consisting of an alpha and a beta subunit. The alpha subunit alone binds IL-3 with low affinity. The beta subunit does not bind IL-3 by itself but is required for the high-affinity binding of IL-3 to the heterodimeric Receptor complex.

SDS-PAGE

